



兔抗 CISH 多克隆抗体

中文名称: 兔抗 CISH 多克隆抗体

英文名称: Anti-CISH rabbit polyclonal antibody

别 名: CIS; G18; SOCS; CIS-1; BACTS2

抗 原: CISH

相关类别: 一抗

储 存: 冷冻 (-20°C)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate


克隆类型: rabbit polyclonal

技术规格

Background:

The protein encoded by this gene contains a SH2 domain and a SOCS box domain. The protein thus belongs to the cytokine-induced STAT inhibitor (CIS), also known as suppressor of cytokine signaling (SOCS) or STAT-induced STAT inhibitor (SSI), protein family. CIS family members are known to be cytokine-inducible negative regulators of cytokine signaling. The expression of this gene can be induced by IL2, IL3, GM-CSF and EPO in hematopoietic cells. Proteasome-mediated degradation of this protein has been shown to be involved in the inactivation of the erythropoietin receptor. Multiple transcript variants enc

	oding different isoforms have been found for this gene.
Applications:	ELISA, IHC
Name of antibody:	CISH
Immunogen:	Fusion protein of human CISH
Full name:	cytokine inducible SH2 containing protein
Synonyms:	CIS; G18; SOCS; CIS-1; BACTS2
SwissProt:	Q9NSE2
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human esophagus cancer and Human thyroid cancer
IHC Recommend dilution:	100-300



mlbio 海聯生物
Good elisakit producers

